


Nucleic acids Extraction

By


Dr/ Rania Ewida

Ass. Prof of Milk Hygiene

Basic structure of the cell



Prokaryotic cell



Eukaryotic cell

Prokaryotic cell

Eukaryotic cell

□ Most DNA extraction protocols consist of

□ Cells lysis

□ Precipitation

□ Washing

□ Resuspension


□ Cells lysis

□ Bacteria

Lysozymes

□ (Tears, egg white, milk, mm)

Breaking peptidoglycan
□ cell wall in Bacteria



Alexander Fleming
□ Nobel Prize 1945



DNA Extraction

□ Cells lysis


□ Plant, Fungi

□ Mechanical Force

Liquid nitrogen and grinding , Sonication,
□ grinding

□ Animal cell

□ Mild Detergent



DNA Extraction



□ Cells lysis

SDS(sodium dodecyl sulfate)

- Remove lipids
- Denature proteins

Proteinase K (65 °C)

- Digest protein
- Inactivate DNase
- Remove Histones
- Very active with SDS





DNA Extraction

□ Cells lysis

□ Phenol/Chloroform

- Separate DNA and RNA from other components
- Denature Proteins



DNA Extraction

□ Precipitation

2-3 volumes cold ethanol 95% with
□ high salt concentration






Figure 4.2 Ethanol precipitation



□ Washing

It is “washed” with a 70% ethanol solution to remove salts and other water soluble impurities but not resuspend the DNA.□

Most salts are soluble in 70% ethanol

□ Resuspension

The clean DNA is now resuspended in a buffer or water to ensure stability and long term storage.

The most commonly used buffer for resuspension is called **1xTE** or **water**






Spin columns



PCR reaction product or
Enzymatic reaction product (**FAPC Buffer**)



Pure DNA fragment

DNA Extraction

PCR from bacteria

Freshly cultured
□ bacterial colonies



1 colony in 100µl water, □
95°C for 5 minutes. □

□ 1-3 µl for PCR directly □ centrifugation



low quality DNA but good
□ enough for routine analyses

DNA Extraction

Direct PCR from blood, cells, tissues and plants

- **Whole Blood:**

use 1 μl directly in 50 μl reaction
or preheat larger volumes 95 °C 15 min (McCusker *et al.*, 1992) (*Even with anticoagulant*)

- **Cells Resuspend :**

10ul of cell culture in water heat 100 °C in PCR machine for 5 minutes use 1-2 μl for PCR

- **Tissues:**


(100 % formamide, heat 95 and 72°C 30 times prior to PCR. Use 2-3 μl for reaction) (Panaccio *et al.*, 1993)

- **Plant (seed):**

(Drilling out a sample from the seed, adding NaOH, heating in a microwave oven and neutralizing with Tris-HCl. (Von Post *et al.*, 2003). Use 1-3 μl for reaction



Overview of DNA Extraction



**Break down
the cell wall &
membranes**



**Centrifuge to
separate the
solids from the
dissolved DNA**



**Precipitate
the DNA
using
alcohol**



**Centrifuge
to separate
the DNA
from the
dissolved
salts and
sugars**



**Dissolve
DNA**



**Wash the DNA
pellet with
Ethanol and dry
the pellet**

Evaluation of Nucleic Acids


- **Spectrophotometrically**

- quantity
- quality



- **Fluorescent dyes**

- gel electrophoresis



DNA quality and concentration



DNA	A_{260}	$1.0 \approx 50 \mu\text{g/ml}$
	A_{260}/A_{280}	$1.6 - 1.8$
RNA	A_{260}	$1.0 \approx 40 \mu\text{g/ml}$
	A_{260}/A_{280}	~ 2.0



DNA Extraction

Analyzing DNA samples

By using gel electrophoresis

Analysis of samples:


Barley (A): This sample is fine

Corn (B): This sample is fine

Oat (C) : This sample is fine

Rice (D) : This sample is fine

Wheat (E): This sample has severe degradation, can work for PCR but should re-extract



DNA Extraction



Thank You!