

1. Which of the following is not an advantage of marsupialization?
 - A. Exposure of very little bone
 - B. Preservation of vital structure
 - C. Rapid healing
 - D. Conserve surgical sutures

2. An empty cavity in the mandible with no lining is most likely to be :
 - A. Aneurysmal bone cyst
 - B. Idiopathic bone cavity
 - C. Dentigerous cyst
 - D. Keratocyst

3. The cyst is deroofed and the surrounding periosteum is sutured to the margins of the cyst wall in:
 - A. Decortication
 - B. Marsupialization
 - C. Saucerization
 - D. Enucleation

4. Ameloblastoma is best managed by:
 - A. Chemotherapy
 - B. Radiotherapy
 - C. Gene Therapy
 - D. Surgical excision

5. The most appropriate method to differentiate between a dentigerous cyst and an ameloblastoma is through:
 - A. Radiographic examination
 - B. Aspiration cytology
 - C. Microscopic examination
 - D. Clinical features

6. When treated with simple curettage, which odontogenic tumor is most likely to recur?

- A. Complex odontoma
- B. Compound odontoma
- C. Odontogenic myxoma
- D. Ameloblastic fibroma

7. Odontoma is treated by:

- A. Excision
- B. Resection
- C. Curettage
- D. Radiotherapy

8. Treatment of ameloblastoma:

- A. Excision of tumor
- B. Resection of the mandible along with the tumor
- C. Incisional biopsy and marsupialization
- D. No active treatment is necessary

9. Osteomyelitis of the jaw can be cured by:

- A. Resection
- B. Physiotherapy
- C. Sequestrectomy with antibiotic treatment
- D. Drainage

10. During development of the teeth, Ectodermal tissues will give rise to

- enamel.

- Dentin
- Pulp
- cementum