

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/384993100>

Sperm rheotaxis as a parameter for laboratory Evaluation of frozen semen in bull

Article · October 2024

CITATIONS

0

READS

22

4 authors, including:



[Haitham Mofadel](#)

Assiut University

4 PUBLICATIONS 0 CITATIONS

[SEE PROFILE](#)



[Hassan Hussein](#)

Assiut University

105 PUBLICATIONS 721 CITATIONS

[SEE PROFILE](#)



[Ahmed M. R. Fath El Bab](#)

Egypt-Japan University of Science and Technology

149 PUBLICATIONS 1,324 CITATIONS

[SEE PROFILE](#)

Sperm rheotaxis as a parameter for laboratory Evaluation of frozen semen in bull

Haitham A. Mofadel^{1*}, Hassan A. Hussein¹, Ahmed M.R. Fath El-Bab², Taymour M. El-Sherry¹

¹Department of Theriogenology, Faculty of Veterinary Medicine, Assiut University, Assiut 71526, Egypt.

²Department of Mechatronics and Robotics, Egypt-Japan University of Science and Technology (E-JUST), Alexandria 21934, Egypt.

ARTICLE INFO

Received: 02 January 2024

Accepted: 19 May 2024

*Correspondence:

Corresponding author: Haitham A. Mofadel
E-mail address: Haitham_Mofadel@aun.edu.eg

Keywords:

Sperm rheotaxis
Cryopreserved semen
Sperm kinematics
Semen evaluation
Cattle.

ABSTRACT

Sperm rheotaxis is a sperm guidance mechanism that helps to guide and select high-quality sperm cells within the female reproductive tract. Semen is usually evaluated using either the traditional subjective method of evaluation or computer assisted sperm analysis (CASA). These methods provide information about sperm concentration, viability and motility parameters and ignoring sperm ability to travel inside the female reproductive tract. Our goal was to detect the individual variations between different sires through evaluation of post thawed frozen semen using sperm rheotaxis. We studied sperm rheotaxis and sperm kinematics inside microfluidic platforms together with subjective semen analysis (motility, viability, and sperm morphology) in post-thawed bull frozen semen in different sires. The results showed that the positive rheotaxis (PR) was higher in the post-thawed frozen semen of some bulls ($P < 0.05$) than in others. Our results also revealed that the bulls with the higher PR% have significantly higher straight-line velocity (VSL) and subjective analysis parameters (motility and viability) ($P < 0.05$) than bull with the lower PR. Furthermore, PR was positively correlated with VSL ($r = 0.67$, $P < 0.0001$) and with subjective motility and viability ($r = 0.57$, $P = 0.007$ and $r = 0.49$; $P = 0.02$, respectively). Thus, we concluded that sperm rheotaxis should be used as a new parameter in evaluation of post-thawed frozen semen quality in bull.

Introduction

Bull fertility is an important factor in cattle reproduction especially after introduction of artificial insemination (AI) technology, as one bull could be used to inseminate large number of cows. However, cryopreservation extends the storage of sperm, associated cryoinjury may cause loss of sperm motility and cause acrosomal damage, mitochondrial membrane depolarization, cryo-capacitation and change in plasma membrane permeability (Bailey *et al.*, 2000). These factors make spermatozoa loss its ability to interact with the female reproductive tract and thus fertilization failure (Chen *et al.*, 2015). Post thawing cryo-survival rate of spermatozoa is considered one of the most vital criteria in bull selection inside AI centers. As it was found that semen of some bulls is more sensitive to cryopreservation than others (Holt, 2000). Male fertility is mainly based on post-thawed semen assessment using conventional parameters such as sperm motility, morphology, viability, biochemical estimations of enzyme release, membrane, and acrosomal integrity. Amongst these parameters motility is believed to be the most important trait associated with spermatozoa fertilizing ability. Motility remains the parameter of choice to reveal the degree of sperm damage caused by the cryopreservation process. Sperm motility evaluation is commonly performed by visual assessment using a microscope equipped with phase contrast optics. Nevertheless, this technique still poor predictor for fertility potential because of its subjectivity (Du Plessis *et al.*, 1988; Fitzpatrick *et al.*, 2002; Holroyd *et al.*, 2002). On the other hand, CASA gives a wider range of spermatozoa motility parameters which in turn offer valuable information about spermatozoa physiological status and therefore has the ability for more accurate prediction of fertility than the parameters evaluated by subjective microscopical semen examination (Christensen *et al.*, 1999; Farrell *et al.*, 1998). Unfortunately CASA parameters give no idea about

the role of selection carried out by the female reproductive tract (Wang and Swerdloff, 2014). Sperm positive rheotaxis (PR) referred to the ability of the spermatozoa to swim against the flow (El-Sherry *et al.*, 2014). Selected spermatozoa based on rheotaxis found to have a good quality parameters such as higher motility (Sarbandi *et al.*, 2021), improved chromatin maturity (De Martin *et al.*, 2017), improved morphology, viability and lower DNA fragmentation (Romero-Aguirregomezcorra *et al.*, 2021). Moreover pre-freezing sperm selection based on rheotaxis showed improvement in cryo-survival rate, post thawed sperm lifespan and in-vivo fertility (Nagata *et al.*, 2019b). In ram, It was reported that high PR resulted in a higher pregnancy rate and lower pregnancy loss and found to be positively correlated with fertility (Abdel-Ghani *et al.*, 2020). Such evidence highlights the importance of sperm rheotaxis in the fertilization process. Thus, this study was designed to detect the individual variation between different cattle sires in post-thawed frozen semen quality using sperm rheotaxis along with CASA parameters for positive rheotactic sperm population and subjective semen analysis.

Materials and methods

Semen sample

Frozen semen from three fertile bulls (21 straws 7 for each). Straws were obtained from three different ejaculates for each bull. Straws were purchased from the Directorate of Veterinary Medicine of the Assiut governorate. For semen quality evaluation (PR%, CASA, and subjective semen analysis) semen straws were thawed in a water bath at 37°C for 30s and assisted directly without any other treatment or incubation except for dilution with sodium citrate.

Post-thaw subjective evaluation of motility, viability, and morphology.

Sperm motility was performed using light microscope. Semen drop was placed on a warm dry clean slide, cover slide was placed over the drop and examined under high power x40. Only sperms with forward progressive motility towards the head are considered to be normal.

The viability of the sperm cells was assessed using live/dead stain as follows: One drop of semen was added to two drops of eosin stain 2% and four drops of Nigrosin stain 10%. Then thin smears are made from this mixture, dried in air and examined using light microscope under high power x40. The percentage of live sperm was calculated by counting two hundred sperm cells in different fields.

Sperm morphology was evaluated using alkaline methyl violet stain according to (Hackett and Macpherson, 1965)

Sperm rheotaxis and kinematics were analyzed using CASA.

The thawed semen straws (n=21) were diluted with sodium citrate in a 1:3 ratios, respectively. As glycerol used in semen cryopreservation prevents flow generation inside microchannel. Then a straight microchannel with 200 µm width, 100 µm height was used to study sperm rheotaxis (PR) and sperm kinematics. The sperm concentration after dilution was around 15×10^6 sperm/ml

PR and kinetic parameters of sperm cells (n =75137 for all examined straws) were determined through a home-made computer-assisted sperm analysis (CASA) system (Department of Mechanical Engineering, Faculty of Engineering, Assiut University, Egypt; the plugin can be downloaded from the following URL: <http://www.assiutmicrofluidics.com/research/casa>) (Elsayed et al., 2015). Sperm cell videos were taken with an Optika XDS-3 inverted microscope with phase contrast (also at 40× objectives) coupled to a Tucsen ISH1000 camera at 30 frames per second. The recorded videos were processed using a home-developed CASA and the following parameters were determined: velocity parameters which included straight-line velocity (VSL, µm/s), average path velocity (VAP, µm/s) and curvilinear velocity (VCL, µm/s) and progression parameters (linearity (LIN=VSL/VCL) and beat cross frequency (BCF, Hz).

Microfluidic Device fabrication

Lithography Chip fabrication

The chip involves 2 PMMA parts, the lower part had the engraved channel structure, and the upper part held inlet ports, as shown in Fig. 1a.

The used channel was made by direct write laser machining technique. VLS3.5 UNEVERSA LASER SYSTEMS with a 30-Watt CO₂ laser tube and 100µm laser beam was used for channel fabrication. We got the best engraving by adjusting the engraving speed to 25mm s⁻¹ (10%) laser head translation speed and laser beam power to 5-Watt (6%) laser beam power to get less roughness at the lowest available dimensions. The channel profile is Gaussian shape as shown in Fig. 1b.

Bonding of the upper and bottom parts (which contained the channel) was done by the thermo-compression method with acetic acid at 115°C and 1 N for 7 min. By heating with acetic acid, better bonding at lower temperature was achieved as well as bonding time. The final dimensions of the microchannel used in this study was 200×100 µm (Width x Height) (Nasser et al., 2019). Microfluidic device used in the present study was shown in Fig. 1c.

Flow generation

Hydrostatic pressure was used to induce liquid flow inside the micro-channel by keeping the liquid level in the inlet reservoir higher than that in the output reservoir at a different height Δh. Hydrostatic flow generation is a simple and low-cost method of generating flow inside micro-

channels and does not suffer from the pulsating flow typical to syringe pumps (Moscovici et al., 2010) Average velocity inside the microchannel was calculated by using the Darcy–Weisbach equation (Munson et al., 2002).

$$(1) \quad V_{av} = \frac{(2\rho g D_h \Delta h)}{C \mu L}$$

where ρ is the density of the liquid, g is the gravitational acceleration, D_h is the hydraulic diameter, Δh is the height difference between the reservoirs, C = friction factor (f) × Reynolds number (Re), μ is the viscosity of the liquid and L is the microchannel length (Munson et al., 2002). The velocity profile inside the channel was calculated using equation (2) for channels with an aspect ratio less than 0.5 (SHAH and AL, 1978).

$$(2) \quad \frac{V}{V_{av}} = \left(\frac{m+1}{m}\right) \left(\frac{n+1}{n}\right) \left[1 - \left(\frac{y}{b}\right)^n\right] \left[1 - \left(\frac{z}{a}\right)^m\right]$$

Where V is the liquid velocity at any location in channel a, and b is the channel width and height, respectively. However, y and z are the coordinates (measured from the centerline) of any point in the channel where V is required, and m and n are numerical parameters dependent on the channel aspect ratio α = b/a according to equations (3) and (4).

$$(3) \quad m = 1.7 + 0.5\alpha^{-1.4}$$

$$(4) \quad n=2 \cdot 0.3(\alpha - 1/3) \begin{cases} \alpha < \frac{1}{3} \\ \alpha \geq \frac{1}{3} \end{cases}$$

Statistics

Data from subjective sperm analyses, sperm rheotaxis, and all sperm kinematic parameters were expressed as mean ± SEM. All data were analyzed using one way analysis of variance (ANOVA) followed by Tukey multiple comparison test to determine the differences between the means of different groups. Linear Relationship has been used to analyze the correlation between VSL, subjective sperm motility and viability with PR. Statistical analysis was performed using Graph Pad Prism version 8.0.0 for Windows, Graph Pad Software, San Diego, California, USA, www.graphpad.com




Fig. 1. (a) Chip schematic drawing. (b) Channel profile. (c) Picture of the microfluidic device used in the present study.

Results

Subjective semen analysis

The results showed that the individual motility and viability of bulls 1 and 3 were significantly higher than those of bull 2 (p <0.0001; Table 1). On the other hand, Results revealed no significant difference among bulls

in sperm morphology (Table 1).

Table 1. Subjective analysis (individual motility, viability, and morphology %) in post-thawed frozen semen.

	Individual motility (%)	Viability (%)	Normal morphology (%)
Bull 1	54.5±1.2 ^a	57.9±0.9 ^a	93.00±0.3
Bull 2	40.0±2.0 ^b	43.5±1.4 ^b	93.50±6.0
Bull 3	50.0±1.5 ^a	54.6±1.0 ^a	92.8±0.7
P value	<0.0001	<0.0001	0.7

Data represented in mean ± SEM. Values with different superscripts are significantly different in column (P < 0.05).

Sperm rheotaxis

Positive rheotaxis was significantly higher in bulls 1 and 3 than in bull 2 (p<0.0001; Table 2).

Sperm kinematics

Two sets of parameters were used. Velocity parameters, which include curvilinear velocity (VCL), straight-line velocity (VSL), and average path velocity (VAP). The other set is progression parameters, which include linearity (LIN=VSL/VCL) and beat / cross frequency (BCF). The VSL was significantly higher in bulls 1 and 3 than in bull 2. The VCL, the VAP and linearity values of bull 1 and 3 were higher than those of bull 2 but without significant difference. The BCF was significantly higher in bull 1 and 2 than in bull 3 (Table 2).

Relationship between VSL, motility and viability with PR.

Our results showed a significant positive correlation between VSL, sperm subjective motility and viability with PR% (r = 0.67 P = <0.0001) (r = 0.57, P = 0.007) (r = 0.49, P = 0.02) (Fig. 2 a, b, c), respectively.




Fig. 2. Showing the relation between (a) VSL with positive rheotaxis (b) motility with positive rheotaxis (PR) (c) Viability with positive rheotaxis (PR). indicate Significant at p < 0.05.

Table 2. The positive rheotaxis % (PR) and sperm kinematics in post-thawed frozen semen.

	N	PR (%)	VCL (μm/s)	VAP (μm/s)	VSL (μm/s)	LIN (VSL/VCL)	BCF (Hz)
Bull 1	31791	54 ± 1.8 ^a	22.7± 0.14	22.6±0.1	17.7±0.04 ^a	0.78 ±0.01	1.7±0.02 ^a
Bull 2	34977	41 ± 1.8 ^b	22.6± 0.09	22.5±0.1	17.4±0.1 ^b	0.77 ±0.004	1.7±0.02 ^a
Bull 3	8369	56 ± 3.9 ^a	23± 0.2	23± 0.2	18±0.1 ^a	0.79 ±0.01	1.6±0.04 ^b
P value		<0.0001	0.4	0.1	<0.0001	0.3	0.01

Data are represented as mean ± SEM. Values with different superscripts are significantly different in column (P < 0.05). N: Sperm number; PR %: Positive rheotaxis %; VCL: Curvilinear velocity; VSL: Straight line velocity; VAP: Average path velocity; LIN=VSL/VCL: linearity and (BCF) beat/cross frequency.

Discussion

The results showed that subjective motility and viability were significantly higher in bulls 1 and 3 than in bull 2, while there was no significant difference in normal sperm morphology among these bulls. Motility still the most valuable parameter to detect cryo-damage of spermatozoa caused by freezing process (Kathiravan *et al.*, 2011). Furthermore, a significant correlation has been found between spermatozoa motility percentage and pregnancy rates in cattle (Holroyd *et al.*, 1993). A strong correlation was detected between post thaw semen viability and field fertility (56-day non return rate) (Januskauskas *et al.*, 2000). The percent of normal morphology of sperm cells was constantly correlated with the calf output (Fitzpatrick *et al.*, 2002). Also, normal sperm morphology and vitality percentage were significantly correlated with fertilization rates when used in-vitro (Du Plessis *et al.*, 1988). On the other hand, standard semen analysis using microscope with phase contrast optics was reported to not represent an accurate fertility predictor because of the subjectivity of the technique (Du Plessis *et al.*, 1988; Kathiravan *et al.*, 2011). Furthermore, the relationship between subjectively evaluated motility and fertility was found to be contradictory: as some reports demonstrated the presence of a significant relationship between them (Correa *et al.*, 1997; Kjoestad *et al.*, 1993; Wood *et al.*, 1986; Zhang *et al.*, 1998) while others could not prove such a relationship (Andersson *et al.*, 1992; Januskauskas *et al.*, 1999; Söderquist *et al.*, 1991).

In the present study, the PR was significantly higher in bulls 1 and 3 than in bull 2. In addition, we found a positive relationship between PR and VSL and subjective semen analysis parameters (motility and viability). Positive sperm rheotaxis was reported as a valuable parameter in the evaluation of ram semen to assess the reproductive performance. As a high percentage of positive rheotaxis >40% achieve higher pregnancy rates (Abdel-Ghani *et al.*, 2020). Moreover, It was found that positive sperm rheotaxis percentage was strongly correlated with fertility in human and it was suggested that such parameter should be used as a fertility indicator in semen analysis and as a selection method in assisted reproductive technology (El-Sherry *et al.*, 2022). The pre-freezing selection of bovine spermatozoa using rheotaxis had a positive effect on the cryo-survival and fertilization ability of sperm cells. These rheotaxis selected spermatozoa showed no change in structural integrity, viability, and biological function after freezing which related to fertilization capability. In addition, when these pre-freezing selected spermatozoa used in-vivo it results in improvement of pregnancy rate in Japanese Black heifers and Holstein repeat breeders (Nagata *et al.*, 2019a). Selected sperms based on rheotaxis were found to have good quality parameters such as higher motility (Sarbandi *et al.*, 2021), improved chromatin maturity (De Martin *et al.*, 2017), improved morphology, viability, and lower DNA fragmentation (Romero-Aguirregomezcorrea *et al.*, 2021). All these proofs, in addition to our results that show strong relation between VSL, motility and viability with positive rheotaxis, confirms the existence of a positive relationship between sperm rheotaxis and high-quality semen samples.

Regarding sperm kinematics, the results revealed that VSL was significantly higher in bulls 1 and 3 than in bull 2. Even the rest of velocity parameters (VCL and VAP) in addition to linearity were insignificantly higher in bulls 1 and 3 than bull 2. The BCF was found to be higher in bulls 1 and 2 than in bull 3. Motion kinetics are very important factors in fertility. As these parameters play a vital role in sperm transportation towards the fertilization site and even when sperm come in contact with the oocyte (Rahman *et al.*, 2013). Furthermore, a strong correlation between motility characteristics (VCL, VSL, LIN) and competitive fertility index had been observed in post-thawed frozen bull semen (AMANN, 1989). A high correlation has been reported between combined motion kinematics and bull fertility with good predictive value (r² = 0.63–0.98) (Farrell *et al.*, 1998). Oliveira *et al.* (2013) have reported a high correlation between CASA parameters combination and bull fertility. Furthermore, it was found that velocity parameters (VCL, VAP, and VSL) of the post-thawed bull semen have a significant correlation with the fertility of the bull (Nagy *et al.*, 2015).

In the present study VSL (progressive velocity) was significantly lower in one bull than in the others. Even the rest of velocity parameters (VCL

and VAP) was lower in the same bull but without significance and This can be explained by the fact that some bull's semen is more sensitive to the freezing process (low freezability) than others (Holt, 2000). The reason behind the decreased motility during cryopreservation is due to the reduction in mitochondrial activity (Yoon *et al.*, 2015). And motility is strongly related to mitochondrial activity (De Lamirande and GAGNON, 1992; Sariözkán *et al.*, 2009). Cryo-damage decreases sperm flagellar activity due to the degenerative process that causes impairment of mitochondrial function (Said *et al.*, 2010). Among all sperm kinematics sperm progressive motility is believed to be vital for mammalian sperm cells to travel through the uterotubal junction and enter the fallopian tube, while non-progressive swimming seems to make it hard for sperm to reach the fertilization site (Gaddum-Rosse, 1981; Shalgi *et al.*, 1992). Moreover, boar spermatozoa isolated from progressive subpopulation showed significant increase in velocity and linearity during in-vitro capacitation. While those who have non-progressive sperms did not practically change their motile parameters (Ramíó *et al.*, 2008). BCF represents the frequency with which the actual trajectory of the sperm crosses the average path trajectory. In a study between high and low fertility Holstein bulls there was a positive correlation between velocity parameters (VCL, VAP, and VSL) and non-return rate. While there was a negative correlation between BCF and non-return rate in post thawed frozen semen (Shojaei *et al.*, 2012). On the other hand BCF was found to have almost no correlation with bull fertility in many studies (Kathiravan *et al.*, 2011). It was reported that the best progression of bull sperms through fluid current take place with high velocity parameters (VCL, VSL and VAP) and low BCF (El-sherry *et al.*, 2017).

From these results, we can conclude that bull 2 has the lowest PR%, subjective motility, and viability among the 3 bulls. Moreover, the sperms of bull 2 appears to swim in a more oscillatory and less progressive pattern than the sperm cells of bulls 1 and 3 as indicated by their lower VSL (progressive velocity). This variation in subjective analysis and sperm kinematics was probably caused by cryo-damage that appears to occur in higher degrees in some bulls that has low freeze ability (bull 2) than others (bull 1,3). Even more, this bull 1 with the low freeze ability had a lower PR % than other bulls.

Conclusion

Bulls with low PR during semen evaluation have low motility and viability and unsatisfactory sperm kinematics in semen evaluation. Therefore, we strongly recommend using sperm rheotaxis as an indicator for semen quality in post-thawed frozen semen in bull.

Conflict of interest

The authors declare that there is no conflict of interest.

References

- Abdel-Ghani, M.A., El-sherry, T., Mahmoud, G., Nagano, M., 2020. Implications of ram sperm rheotaxis analysed by microfluidics for fertility. *Reproduction in Domestic Animals* 55, 1541-1547. <https://doi.org/10.1111/rda.13804>
- AMANN, R.P., 1989. Can the fertility potential of a seminal sample be predicted accurately? *Journal of andrology* 10, 89-98. <https://doi.org/10.1002/j.1939-4640.1989.tb00066.x>
- Andersson, M., Hellman, T., Holmström, B.-G., Jokinen, L., 1992. Computerized and subjective assessments of post-thaw motility of semen from Finnish Ayrshire AI bulls in relation to non-return rates. *Acta Veterinaria Scandinavica* 33, 89-93. [10.1186/BF03546939](https://doi.org/10.1186/BF03546939)
- Bailey, J.L., Bilodeau, J., Cormier, N., 2000. Semen cryopreservation in domestic animals: a damaging and capacitating phenomenon. *Journal of andrology* 21, 1-7. <https://doi.org/10.1002/j.1939-4640.2000.tb03268.x>
- Christensen, P., Brockhoff, P., Lehn-Jensen, H., 1999. The relationship between semen quality and the nonreturn rate of bulls. *Reproduction in domestic animals* 34, 503-507. <https://doi.org/10.1111/j.1439-0531.1999.tb01410.x>
- Correa, J., Pace, M., Zavos, P., 1997. Relationships among frozen-thawed sperm characteristics assessed via the routine semen analysis, sperm functional tests and fertility of bulls in an artificial insemination program. *Theriogenology* 48, 721-731. [https://doi.org/10.1016/S0093-691X\(97\)00296-3](https://doi.org/10.1016/S0093-691X(97)00296-3)
- De Lamirande, E., GAGNON, C., 1992. Reactive oxygen species and human spermatozoa: I. Effects on the motility of intact spermatozoa and on sperm axonemes. *Journal of andrology* 13, 368-378. <https://doi.org/10.1002/j.1939-4640.1992.tb03327.x>
- De Martin, H., Cocuzzo, M.S., Tiseo, B.C., Wood, G.J., Miranda, E.P., Monteleone, P.A., Soares, J.M., Serafini, P.C., Srougi, M., Baracat, E.C., 2017. Positive rheotaxis extended drop: a one-step procedure to select and recover sperm with mature chromatin for intracytoplasmic sperm injection. *Journal of assisted reproduction and genetics* 34, 1699-1708. [10.1007/s10815-017-1024-1](https://doi.org/10.1007/s10815-017-1024-1)
- Du Plessis, Y.P., Nayudu, P.L., Johnston, W.I.H., Baker, H.G., 1988. The use of in vitro fertilization to evaluate putative tests of human sperm function. *Fertility and sterility* 49, 272-277. [https://doi.org/10.1016/S0015-0282\(16\)59715-5](https://doi.org/10.1016/S0015-0282(16)59715-5)
- El-Sherry, T.M., Abdel-Ghani, M.A., Abdel Hafez, H.K., Abdelgawad, M., 2022. Rheotaxis of sperm in fertile and infertile men. *Systems Biology in Reproductive Medicine*, 1-7. <https://doi.org/10.1080/19396368.2022.2141154>
- El-Sherry, T.M., Elsayed, M., Abdelhafez, H.K., Abdelgawad, M., 2014. Characterization of rheotaxis of bull sperm using microfluidics. *Integrative Biology* 6, 1111-1121. <https://doi.org/10.1039/c4ib00196f>
- El-sherry, T., Abdel-Ghani, M., Abou-Khalil, N., Elsayed, M., Abdelgawad, M., 2017. Effect of pH on rheotaxis of bull sperm using microfluidics. *Reproduction in Domestic Animals* 52, 781-790. <https://doi.org/10.1111/rda.12979>
- Elsayed, M., El-Sherry, T.M., Abdelgawad, M., 2015. Development of computer-assisted sperm analysis plugin for analyzing sperm motion in microfluidic environments using Image-J. *Theriogenology* 84, 1367-1377. <https://doi.org/10.1016/j.theriogenology.2015.07.021>
- Farrell, P., Presicce, G., Brockett, C., Foote, R., 1998. Quantification of bull sperm characteristics measured by computer-assisted sperm analysis (CASA) and the relationship to fertility. *Theriogenology* 49, 871-879. [https://doi.org/10.1016/S0093-691X\(98\)00036-3](https://doi.org/10.1016/S0093-691X(98)00036-3)
- Fitzpatrick, L., Fordyce, G., McGowan, M., Bertram, J., Doogan, V., De Faveri, J., Miller, R., Holroyd, R., 2002. Bull selection and use in northern Australia: Part 2. Semen traits. *Animal reproduction science* 71, 39-49. [https://doi.org/10.1016/S0378-4320\(02\)00024-6](https://doi.org/10.1016/S0378-4320(02)00024-6)
- Gaddum-Rosse, P., 1981. Some observations on sperm transport through the uterotubal junction of the rat. *American Journal of Anatomy* 160, 333-341. <https://doi.org/10.1002/aja.1001600309>
- Hackett, A., Macpherson, J., 1965. Some staining procedures for spermatozoa. *The Canadian Veterinary Journal* 6, 55.
- Holroyd, R., Doogan, V., De Faveri, J., Fordyce, G., McGowan, M., Bertram, J., Vankan, D., Fitzpatrick, L., Jayawardhana, G., Miller, R., 2002. Bull selection and use in northern Australia: 4. Calf output and predictors of fertility of bulls in multiple-sire herds. *Animal Reproduction Science* 71, 67-79. [https://doi.org/10.1016/S0378-4320\(02\)00026-X](https://doi.org/10.1016/S0378-4320(02)00026-X)
- Holroyd, R., Entwistle, K., O'Rourke, P., Fordyce, G., 1993. Reproductive traits and mating outcomes of Bos indicus bulls in northern Australia. In: *Proceedings of the VII World Conference on Animal Production*, Edmonton, Canada, pp. 44-45.
- Holt, W., 2000. Fundamental aspects of sperm cryobiology: the importance of species and individual differences. *Theriogenology* 53, 47-58. [https://doi.org/10.1016/S0093-691X\(99\)00239-3](https://doi.org/10.1016/S0093-691X(99)00239-3)
- Januskauskas, A., Gil, J., Söderquist, L., Hrd, M., Hrd, M.C., Johannisson, A., Rodriguez-Martinez, H., 1999. Effect of cooling rates on post-thaw sperm motility, membrane integrity, capacitation status and fertility of dairy bull semen used for artificial insemination in Sweden. *Theriogenology* 52, 641-658. [https://doi.org/10.1016/S0093-691X\(99\)00159-4](https://doi.org/10.1016/S0093-691X(99)00159-4)
- Januskauskas, A., Johannisson, A., Söderquist, L., Rodriguez-Martinez, H., 2000. Assessment of sperm characteristics post-thaw and response to calcium ionophore in relation to fertility in Swedish dairy AI bulls. *Theriogenology* 53, 859-875. [https://doi.org/10.1016/S0093-691X\(00\)00235-1](https://doi.org/10.1016/S0093-691X(00)00235-1)
- Kathiravan, P., Kalatharan, J., Karthikeya, G., Rengarajan, K., Kadirvel, G., 2011. Objective sperm motion analysis to assess dairy bull fertility using computer-aided system-a review. *Reproduction in Domestic Animals* 46, 165-172. <https://doi.org/10.1111/j.1439-0531.2010.01603.x>
- Kjoestad, H., Ropstad, E., Andersen Berg, K., 1993. Evaluation of spermatological parameters used to predict the fertility of frozen bull semen. *Acta Veterinaria Scandinavica* 34, 299-303. <https://doi.org/10.1186/bf03548194>
- Moscovici, M., Chien, W.-Y., Abdelgawad, M., Sun, Y., 2010. Electrical power free, low dead volume, pressure-driven pumping for microfluidic applications. *Biomicrofluidics* 4, 046501. <https://doi.org/10.1063/1.3499939>
- Munson, B., Young, D., Okishi, T., 2002. *Fundamentals of Fluids Mechanics*, Edisi 4, Jhon Wiley & Sons, Inc, New York.
- Nagata, M.B., Egashira, J., Katafuchi, N., Endo, K., Ogata, K., Yamanaka, K., Yamanouchi, T., Matsuda, H., Hashiyada, Y., Yamashita, K., 2019a. Bovine sperm selection procedure prior to cryopreservation for improvement of post-thawed semen quality and fertility. *Journal of animal science and biotechnology* 10, 1-14. [10.1186/s40104-019-0395-9](https://doi.org/10.1186/s40104-019-0395-9)
- Nagata, M.B., Egashira, J., Katafuchi, N., Endo, K., Ogata, K., Yamanaka, K., Yamanouchi, T., Matsuda, H., Hashiyada, Y., Yamashita, K., 2019b. Bovine sperm selection procedure prior to cryopreservation for improvement of post-thawed semen quality and fertility. *Journal of animal science and biotechnology* 10, 1-14. [10.1186/s40104-019-0395-9](https://doi.org/10.1186/s40104-019-0395-9)
- Nagy, Á., Polichronopoulos, T., Gáspárdy, A., Solti, L., Cseh, S., 2015. Correlation between bull fertility and sperm cell velocity parameters generated by computer-assisted semen analysis. *Acta Veterinaria Hungarica* 63, 370-381. [10.1556/004.2015.035](https://doi.org/10.1556/004.2015.035)
- Nasser, G.A., Fath El-Bab, A.M., Abdel-Mawgood, A.L., Mohamed, H., Saleh, A.M., 2019. CO₂ laser fabrication of PMMA microfluidic double T-junction device with modified inlet-angle for cost-effective PCR application. *Micromachines* 10, 678. [10.3390/mi10100678](https://doi.org/10.3390/mi10100678)
- Oliveira, L.Z., de Arruda, R.P., de Andrade, A.F.C., Celeghini, E.C.C., Reeb, P.D., Martins, J.P.N., dos Santos, R.M., Beletti, M.E., Peres, R.F.G., Monteiro, F.M., 2013. Assessment of in vitro sperm characteristics and their importance in the prediction of conception rate in a bovine timed-AI program. *Animal reproduction science* 137, 145-155. [10.1016/j.anireprosci.2013.01.010](https://doi.org/10.1016/j.anireprosci.2013.01.010)
- Rahman, M.S., Lee, J.-S., Kwon, W.-S., Pang, M.-G., 2013. Sperm proteomics: road to male fertility and contraception. *International journal of endocrinology* 2013, 10.1155/2013/360986
- Ramíó, L., Rivera, M., Ramírez, A., Concha, I., Peña, A., Rigau, T., Rodríguez-Gil, J., 2008. Dynamics of motile-sperm subpopulation structure in boar ejaculates subjected to "in vitro" capacitation and further "in vitro" acrosome reaction. *Theriogenology* 69, 501-512. [10.1016/j.theriogenology.2007.10.021](https://doi.org/10.1016/j.theriogenology.2007.10.021)
- Romero-Aguirregomez-cortá, J., Laguna-Barraza, R., Fernández-González, R., Štiavníká, M., Ward, F., Cloutery, J., McAuliffe, D., Larsen, P.B., Grabrucker, A.M., Gutiérrez-Adán, A., 2021. Sperm selection by rheotaxis improves sperm quality and early embryo development. *Reproduction* 161, 343-352. [10.1530/REP-20-0364](https://doi.org/10.1530/REP-20-0364)
- Said, T.M., Gaglani, A., Agarwal, A., 2010. Implication of apoptosis in sperm cryoinjury. *Reproductive biomedicine online* 21, 456-462. [10.1016/j.rbmo.2010.05.011](https://doi.org/10.1016/j.rbmo.2010.05.011)
- Sarbandi, I.R., Lesani, A., Moghimi Zand, M., Nosrati, R., 2021. Rheotaxis-based sperm separation using a biomimicry microfluidic device. *Scientific Reports* 11, 18327. [10.1038/s41598-021-97602-y](https://doi.org/10.1038/s41598-021-97602-y)
- Sariözkán, S., Bucak, M.N., Tuncer, P.B., Ulutaş, P.A., Bilgen, A., 2009. The influence of cysteine and taurine on microscopic-oxidative stress parameters and fertilizing ability of bull semen following cryopreservation. *Cryobiology* 58, 134-138. [10.1016/j.cryobiol.2008.11.006](https://doi.org/10.1016/j.cryobiol.2008.11.006)
- SHAH, R., AL, L., 1978. LAMINAR FLOW FORCED CONVECTION IN DUCTS. A SOURCE BOOK COMPACT HEAT EXCHANGER ANALYTICAL DATA.
- Shalgi, R., Smith, T.T., Yanagimachi, R., 1992. A quantitative comparison of the passage of capacitated and uncapacitated hamster spermatozoa through the uterotubal junction. *Biology of Reproduction* 46, 419-424. [10.1095/biolreprod46.3.419](https://doi.org/10.1095/biolreprod46.3.419)
- Shojaei, H., Kroetsch, T., Wilde, R., Blondin, P., Kastelic, J., Thundathil, J., 2012. Moribund sperm in frozen-thawed semen, and sperm motion end points post-thaw and post-swim-up, are related to fertility in Holstein AI bulls. *Theriogenology* 77, 940-951. [10.1016/j.theriogenology.2011.09.026](https://doi.org/10.1016/j.theriogenology.2011.09.026)
- Söderquist, L., Rodríguez-Martinez, H., Janson, L., 1991. Post-thaw motility, ATP content and cytochrome C oxidase activity of AI bull spermatozoa in relation to fertility. *Journal of Veterinary Medicine Series A* 38, 165-174. [10.1111/j.1439-0442.1991.tb00998.x](https://doi.org/10.1111/j.1439-0442.1991.tb00998.x)
- Wang, C., Swerdloff, R.S., 2014. Limitations of semen analysis as a test of male fertility and anticipated needs from newer tests. *Fertility and sterility* 102, 1502-1507. [10.1016/j.fertnstert.2014.10.021](https://doi.org/10.1016/j.fertnstert.2014.10.021)
- Wood, P., Foulkes, J., Shaw, R., Melrose, D., 1986. Semen assessment, fertility and the selection of Hereford bulls for use in AI. *Reproduction* 76, 783-795. [10.1530/jrf.0.0760783](https://doi.org/10.1530/jrf.0.0760783)
- Yoon, S.-J., Kwon, W.-S., Rahman, M.S., Lee, J.-S., Pang, M.-G., 2015. A novel approach to identifying physical markers of cryo-damage in bull spermatozoa. *PLoS One* 10, e0126232. [10.1371/journal.pone.0126232](https://doi.org/10.1371/journal.pone.0126232)
- Zhang, B., Larsson, B., Lundheim, N., Rodríguez-Martinez, H., 1998. Sperm characteristics and zona pellucida binding in relation to field fertility of frozen-thawed semen from dairy AI bulls. *International Journal of Andrology* 21, 207-216. [10.1046/j.1365-2605.1998.00114.x](https://doi.org/10.1046/j.1365-2605.1998.00114.x)